

$$E \frac{d^3 \sigma^{\pi^0}}{dp^3} \text{ [mb / GeV}^2\text{]}$$

■ PHENIX data

# GLOBAL **QCD** ANALYSES & PARTON DENSITIES

Rodolfo Sassot  
Universidad de Buenos Aires

PHENIX SpinFest 2014

— DSS  
- - - KRE  
... AKK  
▨ scale uncertainty



# Second Lecture: Fragmentation Functions

2.1 FFs and jets:

2.2 SIA, SIDIS and pp:


2.3 Global FFs fits: DSS

2.4 Uncertainties: Lagrange

2.5 Uncertainties: Hessian

2.6 DSS-II

2.7 Outlook



$$\sigma = f_i \otimes f_j \otimes \sigma_{ijk} \otimes D_k^\pi$$




## 2.1 FFs and jets

Looking for **FFs**:



$$z \equiv \frac{E_h}{E_q} = \frac{2E_h}{Q}$$

$$\frac{d\sigma}{dz}(e^+e^- \rightarrow h X) = \sum_q \sigma(e^+e^- \rightarrow q\bar{q}) [D_q^h(z) + D_{\bar{q}}^h(z)]$$



$$D_q^h(z) \longrightarrow D_q^h(z, Q^2)$$


- ➡ describe collinear transition of a parton  $i$  into hadron  $h$
  - ➡ universal (process independent)
  - ➡ known scale (energy) dependence
  - ➡ unavoidable if identified hadronic final state
- same theoretical footing as PDFs*



## 2.1 FFs and jets




*why does a collinear framework do so well?*  
*nature prefers collinear emission?*





## 2.1 FFs and jets


A nice example:  $\hat{\sigma}(e^+e^- \rightarrow q\bar{q}g)$



$$z_q \equiv \frac{2E_q}{Q} \quad z_{\bar{q}} \equiv \frac{2E_{\bar{q}}}{Q} \quad z_g \equiv \frac{2E_g}{Q}$$


$$z_q + z_{\bar{q}} + z_g = 2$$

$$\frac{d\hat{\sigma}}{dz_q dz_{\bar{q}}} = \sigma_0 \frac{2\alpha_s}{3\pi} \frac{z_q^2 + z_{\bar{q}}^2}{(1 - z_q)(1 - z_{\bar{q}})}$$







## 2.2 SIA, SIDIS and pp



$$d\sigma = \hat{\sigma}_i \otimes D_i^h$$



$$d\sigma = f_i \otimes \hat{\sigma}_{ij} \otimes D_j^h$$



$$d\sigma = f_i \otimes f_j \otimes \hat{\sigma}_{ijk} \otimes D_k^h$$

$$e^+e^- \rightarrow (\gamma, Z) \rightarrow H$$

clean: only FFs

very precise LEP, Belle, BaBar data

tagged heavy flavors


only information on  $[D_q^H(z, Q^2) + D_{\bar{q}}^H(z, Q^2)]$

“singlet”  $\Sigma \equiv D_u^H + D_{\bar{u}}^H + D_d^H + D_{\bar{d}}^H + D_s^H + D_{\bar{s}}^H + \dots$


suppressed gluon  $\frac{\alpha_s(Q^2)}{2\pi} D_g^H(z, Q^2)$




## 2.2 SIA, SIDIS and pp



$$d\sigma = \hat{\sigma}_i \otimes D_i^h$$



$$d\sigma = f_i \otimes \hat{\sigma}_{ij} \otimes D_j^h$$



$$d\sigma = f_i \otimes f_j \otimes \hat{\sigma}_{ijk} \otimes D_k^h$$

$$l + p \longrightarrow l' + H + X$$

$$F_1^H(x, z, Q^2) = \frac{1}{2} \sum_{q, \bar{q}} e_q^2 f_q(x, Q^2) D_q(z, Q^2)$$


NLO

effective charge: separation


$$+ \frac{\alpha_s}{2\pi} \sum_{q, \bar{q}} e_q^2 [f_q \otimes C_{qq} \otimes D_q^H + f_q \otimes C_{gq} \otimes D_g^H + f_g \otimes C_{qg} \otimes D_q^H]$$




## 2.2 SIA, SIDIS and pp



$$d\sigma = \hat{\sigma}_i \otimes D_i^h$$



$$d\sigma = f_i \otimes \hat{\sigma}_{ij} \otimes D_j^h$$



$$d\sigma = f_i \otimes f_j \otimes \hat{\sigma}_{ijk} \otimes D_k^h$$


$$l + p \longrightarrow l' + H + X$$

charge and flavor separation  
complementary  $z, Q^2$  ranges  
mainly light flavors


**requires precise PDFs!**




## 2.2 SIA, SIDIS and $p\bar{p}$




$$d\sigma = \hat{\sigma}_i \otimes D_i^h$$




$$d\sigma = f_i \otimes \hat{\sigma}_{ij} \otimes D_j^h$$



$$d\sigma = f_i \otimes f_j \otimes \hat{\sigma}_{ijk} \otimes D_k^h$$



$$\sim f_q \otimes f_{\bar{q}} \otimes C_{q\bar{q}g} \otimes D_g^h$$




$$\sim f_g \otimes f_g \otimes C_{ggg} \otimes D_g^h$$


$$p + p \longrightarrow H + X$$




## 2.2 SIA, SIDIS and $p\bar{p}$



$$d\sigma = \hat{\sigma}_i \otimes D_i^h$$



$$d\sigma = f_i \otimes \hat{\sigma}_{ij} \otimes D_j^h$$



$$d\sigma = f_i \otimes f_j \otimes \hat{\sigma}_{ijk} \otimes D_k^h$$

large and direct gluon contribution  
several sub-processes  
RHIC Tevatron LHC benchmarks

$p + p \longrightarrow H + X$   
requires precise PDFs  
“convoluted”  
large TH uncertainty




## 2.2 SIA, SIDIS and $p\bar{p}$

example:  $p\bar{p}@7\text{ TeV}$

$p\text{QCD}$  uncertainties  
> 50%  $p_T < 5\text{ GeV}$

PDFs uncertainties  
5-10%



nevertheless our best grip on gluons...



## 2.2 SIA, SIDIS and $p\bar{p}$

example:  $pp@7\text{ TeV}$


“convoluted”

gluon PDF at low  $p_T$   
but....

gluon FF everywhere

range in  $z$  ( $\langle z \rangle \sim 0.45$ )

range in  $x$  ( $\langle x \rangle \sim 0.01-0.1$ )



at fixed  $p_T$   $\eta$  selects



## 2.2 SIA, SIDIS and pp

multiple convolutions  
time consuming

$$d\sigma = \sum_{abc} \int \int \int f_a f_b d\hat{\sigma}_{ab \rightarrow cX} D_c dx_a dx_b dz_c \quad \frac{1}{2\pi i} \int_{C_n} dn z_c^{-n} D_c^n$$

**Mellin space:**  $D_i^n(\mu^2) = \int_0^1 dz z^{n-1} D_i(z, \mu^2)$

~ O(10<sup>2</sup>) moments accurate representation of FFs (and PDFs)



transforms: convolutions into products (of numbers)  
differential equations into linear equations

*fast and easy  
if known functional form  
(known analytic continuation)*

back to z space (inversion) very fast (complex analysis)

$$d\sigma = \frac{1}{2\pi i} \sum_{abc}$$

$$\int_{C_n} dn$$

Mellin  
inversion

$$D_c^n$$

FIT

$$\int \int \int z_c^{-n} f_a f_b d\hat{\sigma}_{ab \rightarrow cX} dx_a dx_b dz_c$$

Precompute once and save grids



## 2.3 Global FFs fits: DSS

*several FFs parameterizations for  $\pi, K, p$  and  $h^\pm$  ( $\Lambda, \eta, D_c \dots$ )*

CGGRW(1994), BFGW(2000)

Bourhis et al (2001)

BKK(1995), KKP(2000), AKK(2005,2008)  
*include  $pp$*

Albino, Kniehl, Kramer (1995-2008)

KRE(2000)

Kretzer (2000)

HKNS(2007)

Hirai, Kumano, Nagai, Sudoh (2007)

DSS(2007)

*only global analysis*

de Florian, RS, Stratmann (2007)

LSS(2013)  
*new SIDIS only*

Leader, Sidorov, Stamenov (2013)

SLA-only fits: no charge separation or ad-hoc assumption

$$D_{\bar{q}}^{h^+}(z, Q^2) = (1 - z) D_q^{h^+}(z, Q^2)$$



## 2.3 Global FFs fits: DSS

fragmentation functions for  $\pi^+, \pi^-, \pi^0, K^+, K^-, K^0, \bar{K}^0, p, \bar{p}, n, \bar{n}, h^+, h^-$   
$$h^+ = \pi^+ + K^+ + p + res^+$$

LO and NLO global fits available

SIA data includes: TPC, TASSO, SLD, ALEPH, DELPHI, OPAL + “flavor” tag

SIDIS data from : HERMES, EMC(h)  $Q > 1 \text{ GeV}$

pp data from : PHENIX, STAR, BRAHMS, CDF, UAI, UA2  $p_T > 1 \text{ GeV}$  ( $\eta$ )

Estimation of uncertainties using Lagrange multipliers (2007)

Improved Hessian approach (2012)

Data and error estimate update (2014) DSS II



## 2.3 Global FFs fits: DSS

- Flexible parametrization

$$D_i^H(z, Q_0^2) = N_i z^{\alpha_i} (1 - z)^{\beta_i} [1 + \gamma_i (1 - z)^{\delta_i}]$$

at initial scale

$$Q_0^2 = 1 \text{ GeV}^2 \quad u, d, s, g$$

$$Q_0^2 = m_Q^2 \quad c, b$$

with

$\alpha_s$  and  $\Lambda_{QCD}$  from MRST 2002

- Try to avoid Isospin symmetry assumptions

- *allow for possible breaking of SU(3) of sea and SU(2) in favored distributions*

- *unless data can not discriminate for unfavored fragmentations*

$$D_{d+\bar{d}}^{\pi^+} = N D_{u+\bar{u}}^{\pi^+}$$

$$D_s^{\pi^+} = D_{\bar{s}}^{\pi^+} = N' D_{\bar{u}}^{\pi^+}$$

$$D_{\bar{u}}^{\pi^+} = D_d^{\pi^+}$$

$$D_{\bar{u}}^{K^+} = D_s^{K^+} = D_d^{K^+} = D_{\bar{d}}^{K^+}$$

- Normalizations for different experiments (if not included in syst.)






## 2.3 Global FFs fits: DSS

SLA: still works very well  
within the global fit

factorization ✓  
universality ✓

z-cut  $z > 0.05$   
(0.1 for Kaons/Protons)



large errors at  $z > 0.5$



## 2.3 Global FFs fits: DSS


Good description  
of SIDIS multiplicities

ad-hoc charge separation  
from Kretzer fails

large  $z$  covered

*hard to fit!*

Hermes data (not final)






## 2.3 Global FFs fits: DSS

Not so nice for Kaons

ad-hoc charge separation  
from **HKNS** fails  
factor of 2 or 3!!


even harder to fit!





## 2.3 Global FFs fits: DSS


pp data: look impressive (in log scale)






## 2.3 Global FFs fits: DSS

Agreement with data not included in the fit



*STAR charged pions: forgotten!*




*$p_T$ -dependent SIDIS: too difficult!*



## 2.3 Global FFs fits: DSS

similar singlet: SIA

differences in sea:  
explains SIDIS and pp



huge differences in the gluon at large  $z$  : pp




## 2.4 Uncertainties: Lagrange Multipliers

DSS use LM technique to estimate uncertainties (from exp. errors)

**most robust method:** no assumptions on  $\chi^2(\{a_i\})$  or  $\mathcal{O}(\{a_i\})$

*just compute it!*



minimize:

$$\Phi(\lambda_i, \{a_j\}) = \chi^2(\{a_j\}) + \sum_i \lambda_i \mathcal{O}_i(\{a_j\})$$

*See how fit deteriorates when FFs forced to give different prediction for  $\mathcal{O}_i$*

$\Delta\chi^2$  *should be parabolic if data set can determine the observable*

what  $\Delta\chi^2$ ?

*parameter fitting*

68% 90%

1 2.71

*hypothesis testing*

$\sim \sqrt{N}$

*“pragmatic”*

*whatever it takes to have the fitted data within bands*

$$\Delta\chi^2_{68} = \xi_{68} - \xi_{50} \quad \int_0^{\xi_{68}} d\chi^2 P_N(\chi^2) = 0.68$$



## 2.4 Uncertainties: Lagrange Multipliers


example: truncated moments:

$$\int_{0.2}^1 z D_i^H(z, Q = 5 \text{ GeV}) dz$$

$$\Delta\chi^2 = 15 (\sim 2\%)$$

$$u < 5\% \quad s \sim 10\%$$

dominated by  $z \sim 0.2$




factor of 2 larger for Kaons  $\sim 20\text{-}25\%$  for protons



## 2.4 Uncertainties: Lagrange Multipliers

Individual profiles for data subsets: impact & interplay



Constrained parabola as a result of global fit

not user friendly!



## 2.5 Uncertainties: Standard Hessian

$$D_q^h(z, p_{T0}^2) \longrightarrow S \{a_i\} \quad \text{set of fit parameters}$$

$$\text{best fit} \quad S_0 \{a_i^0\} \quad \chi_0^2 = \chi^2(\{a_i^0\})$$

assuming quadratic  
profile

$$\Delta\chi^2 \equiv \chi^2(\{a_i\}) - \chi^2(\{a_i^0\}) \simeq \sum_{ij} H_{ij} y_i y_j$$


$$H_{ij} \equiv \frac{1}{2} \frac{\partial^2 \chi^2}{\partial y_i \partial y_j} \quad y_i \equiv a_i - a_i^0$$

assuming linear  
propagation

$$\Delta\mathcal{O} \simeq \left[ \Delta\chi^2 \sum_{ij} \frac{\partial \mathcal{O}}{\partial y_i} (H^{-1})_{ij} \frac{\partial \mathcal{O}}{\partial y_j} \right]^{1/2}$$



## 2.5 Uncertainties: Improved Hessian



$$\sum_j H_{ij} v_{jk} = \epsilon_k v_{ik}$$

$$a_i - a_i^0 = \sum_k v_{ik} s_k z_k$$

$$\Delta\chi^2 = \sum_k z_k^2$$

$S_k^\pm$  *Hessian eigenvector sets*

$$\Delta\mathcal{O} \simeq \left[ \frac{1}{2} \sum_k [\mathcal{O}(S_k^+) - \mathcal{O}(S_k^-)]^2 \right]^{1/2}$$

*help with correlations/variations  
easier/faster than LM  
replicas for re-weighting*



## 2.6 DSS-II

DSS + M. Epele, R. Hernández Sanchez / 4xx.xxxx

*mostly a DSS fit but with new SIA, SIDIS, RHIC and LHC data (pions)*

analytic normalization

$$\chi^2 = \sum_j \sum_i \left( \frac{\sigma_i^{exp} - \mathcal{N}_j \sigma_i^{th}}{\Delta \sigma_i^{exp}} \right)^2 + \sum_j \left( \frac{1 - \mathcal{N}_j}{\Delta \sigma_j^{exp}} \right)^2$$

flexible parameterizations

$$D_i^{\pi^+}(z, Q_0^2) = N_i z^{\alpha_i} \left[ (1-z)^{\beta_i} + \gamma_i (1-z)^{\delta_i} \right]$$

$$D_{\bar{s}}^{\pi^+}(z, Q_0^2) = N_s z^{\alpha_s} D_u^{\pi^+}(z, Q_0^2)$$

MSTW input PDFs and  $\alpha_s$


pT cuts in pp data

68 and 90 % CL error estimates (improved hessian)




# New SIA

new Belle and BaBar (large  $z$ , lower  $Q$ )



$z \sim 0.05-0.2$   $z > 0.2$


$Q \sim M_Z$



very precise up to  $z \sim 0.8$

$Q \sim 10 \text{ GeV}$






# New SIDIS

final HERMES (p & d) new COMPASS

same kinematics

much better agreement  
(MSTW?)

deuteron~flavor separation






# New SIDIS

final HERMES (p & d) new COMPASS

*bins in  $x$*


*continuous  $z$*

*larger  $Q^2$*





# New SIDIS

(data-theory) / theory







# New pp RHIC and LHC




# New pp RHIC and LHC







FFs:



# Is the Hessian safe?

$$D_i^{\pi^+}(z, Q_0^2) = N_i z^{\alpha_i} [(1-z)^{\beta_i} + \gamma_i (1-z)^{\delta_i}]$$



SU(3):

$$D_s^{\pi^+}(z, Q_0^2) = N_s z^{\alpha_s} D_u^{\pi^+}(z, Q_0^2)$$

SU(2):

$$D_{u+\bar{u}}^{\pi^+}(z, Q_0^2) = N_{d+\bar{d}} D_{u+\bar{u}}^{\pi^+}(z, Q_0^2)$$



# Old and new numbers

DSS

DSS-II

GLOBAL	843.7/392 (2.15)	1154.7/953 (1.21)
LEP-SLAC	500.1/260 (1.92)	412.7/260 (1.58)
BELLE-BABAR	-	90.4/117 (0.77)
HERMES	188.2/64 (2.94)	175.0/128 (1.36)
COMPASS	-	403.2/398 (1.01)
RHIC	160.8/68 (2.36)	43.0/38 (1.13)
LHC	-	27.7/12 (2.31)

# Outlook:

## Experiment

SIDIS: Compass K

SIDIS: Hermes K ✓

SIDIS: Zeus K,  $\Lambda$  ✓

SIA: Belle/BaBar K ✓

pp (K, p,  $\bar{p}$ ,  $\eta$ ,  $\Lambda$ ,  $h^\pm$ ) ratios?

EIC (combined PDFs & FFs?)

## Theory

SIDIS at NNLO

pp NNLO?

heavy quark masses


Threshold resummation

More/different global fits to compare





$$\int_0^{\xi_{68}} d\chi^2 \frac{(\chi^2)^{N/2-1} e^{-\chi^2/2}}{2^{N/2} \Gamma(N/2)} = 0.68$$





# Bibliography

## Introductory:

F. Halzen A. Martin, Quarks and Leptons: Introductory course in Modern Particle Physics, Wiley

T.-P. Cheng, L.-F. Li, Gauge Theory of Elementary Particle Physics, Oxford University Press

## Advanced:

G. Dissertori, I. Knowles, M. Schmelling, Quantum Chromodynamics: High Energy Experiments  
And Theory, Oxford University Press.

R.K. Ellis, W.J. Stirling, B.R. Webber, QCD and Collider Physics, Cambridge University Press,

T. Muta, Foundations of Quantum Chromodynamics, World Scientific

F. J. Yndurain, The Theory of Quark and Gluon Interactions, Springer

G. Altarelli, arXiv:1303.2842

## PDFs:

S. Forte, G. Watt, arXiv:1301.6754

A. de Roeck, R. Thorne, arXiv:1103.0555

## Resource Letter:

A. S. Kronfeld, C. Quigg , arXiv:1002.5032